arXiv:1703.04503v1 [math.AG] 13 Mar 2017

A note on rank 1 log extendable isocrystals on simply connected open varieties

Efstathia Katsigianni*

Abstract

In 2010 de Jong proposed a p-adic version of Gieseker's conjecture: if $\pi_1^{\text{ét}}(X) = 1$, for X smooth connected projective variety, then any isocrystal on X is constant. This was proven by Esnault and Shiho [ES16],[ES15] under some additional assumptions. We show that the conjecture holds in the case of a non-proper variety and for rank 1 log extendable isocrystals.

Introduction

For a complex connected smooth projective variety X one can define the étale fundamental group $\pi_1^{\text{ét}}(X)$, which classifies all the finite étale coverings of X and is isomorphic to the profinite completion of its topological fundamental group. On the other hand we have the category of $\mathcal{O}_X$-coherent $\mathcal{D}_X$-modules, which is equivalent via the Riemann-Hilbert correspondence to the category of finite dimensional representations of the topological fundamental group. By a result of Malčev [Mal40] and Grothendieck [Gro70] we know how this category relates to $\pi_1^{\text{ét}}(X)$: if $\pi_1^{\text{ét}}(X)$ is trivial, then there are no non-constant $\mathcal{O}_X$-coherent $\mathcal{D}_X$-modules.

Naturally, one asks if the same holds true in positive characteristic. This conjecture was formulated by Gieseker [Gie75] and proven by Esnault and Mehta in [EM10].

In 2010 de Jong proposed a p-adic version of this conjecture: if $\pi_1^{\text{ét}}(X) = 1$, then any isocrystal $\mathcal{E} \in \text{Crys}(X/W)_{\mathbb{Q}}$ is constant.

This question was addressed in great detail in [ES16] and in [ES15], where various subcases were considered. There the authors prove, among other results, that for a smooth connected projective variety over an algebraically closed field of positive characteristic, the triviality of the étale fundamental group implies that any convergent isocrystal, which is filtered so that the associated graded is a sum of rank 1 convergent isocrystals, is contant [ES16, Theorem 0.1]. In particular, it is

*Freie Universität Berlin, Arnimallee 3, 14195, Berlin, Germany. Supported by Berlin Mathematical School. Email: katsief@zedat.fu-berlin.de.

also proven that rank 1 isocrystals (not necessarily convergent) are also trivial in this case [ES15, Theorem 1.2].

In the present work we want to extend the aforementioned results of [ES16] to the case of a rank one log extendable isocrystal on a non-proper variety with trivial étale fundamental group. Actually we just need to assume that its abelian quotient is trivial. More precisely, we prove the following:

Theorem 0.1. *Let U be a smooth connected variety over an algebraically closed field k of positive characteristic, that admits a good compactification X, where $X - U = \cup_{i\in I} Z_i =: Z$ is a simple strict normal crossings divisor. If $\pi_1^{ab}(U) = 1$ and $\mathcal{L}$ is a rank one isocrystal on U that extends to a log isocrystal on X, then $\mathcal{L}$ is trivial.*

For the proof of this result we adapt mainly the strategy [ES16, Proposition 3.6] and [ES15, Sections 2.1, 4]. In more detail, starting with a rank one isocrystal $\mathcal{L}$ on U, as in the assumptions of the theorem, we denote by L a locally free lattice of it. First we prove that the log crystalline Chern class $c_1^{logcrys}(L_X)$ of the value of L on X is equal to zero and using this we conclude in Lemma 2.2, that the line bundle L_X to some power N is then of the form $\mathcal{O}_X(\sum a_i Z_i)$ with $\mathbb{Q}$-coefficients. For convenience we denote by E the crystal $L^{\otimes N}$.

Having this information about E_X doesn't allow us though to conclude that E_X is trivial as a log crystal on X/k. However, in Section 3 we adapt the discussion of [ABV05, §6] to our situation and obtain a residue exact sequence:

(0.1)

$$H^1_{crys}(X/W, \mathcal{O}^\times_{X/W}) \to H^1((X,M)/W, \mathcal{O}^\times_{X/W})^{log}_{crys} \to \oplus W(k)[Z_i] \to H^2(X/W, \mathcal{O}^\times_{X/W})$$

It allows us to conclude (see discussion in Section 3), that E is actually "controlled" by its value on X, or more specifically that the map $H^1((X,M)/W, \mathcal{O}^\times)^{log}_{crys} \to H^1(X, \mathcal{O}^\times_X)$ is injective, while one also derives from (0.1) that there is an injection

$$H^1((X,M)/W, \mathcal{O}^\times)^{log}_{crys} \hookrightarrow \oplus W(k)[Z_i]. \tag{0.2}$$

So, one obtains that E on X has actually the form of a sum with $\mathbb{Z}$ coefficients and that E_U is trivial. By Lemma 3.2 we then conclude that one can choose an extension of E to X such that E_X is the trivial crystal on X/k.

In Theorem 4.1 we then use a deformation argument (adapted from [ES16]) to prove that E is the trivial crystal. Finally, we conclude that L and therefore also $\mathcal{L}$ are trivial, using Lemmas 4.2 and 4.3.

As a corollary of this theorem and using the observations made in Section 3, we also obtain in Corollary 4.4 that extensions in $I_{crys}(X/W)^{log}$ of the trivial log isocrystal by itself are constant and so isocrystals on the open variety whose extensions are of this form, are also constant.

Section 5 is actually not necessary for the proof of Theorem 0.1. There we construct the residue map $\oplus W(k)[Z_i] \to H^2(X/W, \mathcal{O}^\times_{X/W})$ appearing in (0.1) assuming that X admits a smooth lift over W.

Finally, we would like to point out the steps of the proof where we use the triviality of $\pi_1^{\mathrm{ab}}(U)$. Precisely, in Lemma 2.2 we use that the maximal pro-l-quotient $\pi_1^{ab,(l)}(U)$ is trivial for $l \neq p$ to obtain that $H^0(U, \mathcal{O}_U^\times) = k^\times$, while $\pi^{\mathrm{ab}}(U) = 1$ also implies $\mathrm{NS}(U) = \mathrm{Pic}(U)$. For the observations of Section 3, the triviality of the abelian quotient $\pi_1^{ab}(U)$ also suffices. Indeed, there we use that $H^1_{crys}(X/W) = H^1_{crys}(X/W, \mathcal{O}_X^\times) = 0$ [ES16, Theorem 0.1.1 and Proposition 2.9] and, as remarked in [Esn16, Theorem 4.3.1], the triviality of the abelian quotient is indeed enough to derive these equalities. The same equalities are also mainly used in Theorem 4.1 together with the fact that $\mathrm{NS}(X) = \mathrm{Pic}(X)$. However, to conclude Section 3 and for Lemma 4.2 we use that Kummer coverings of U are trivial, thus we have to assume that $\pi_1^{\mathrm{tame,ab}}(U) = 1$, since k is algebraically closed.

Acknowledgements: This work is part of my PhD project at the Freie Universität Berlin. I would like to thank my advisor Prof. Hélène Esnault for her guidance and constant support. I would also like to thank Prof. Tomoyuki Abe and Prof. Atsushi Shiho for their most valuable advice and for the very helpful discussions we had during their visits in Berlin.

1 Preliminaries and notation

Let U be a smooth connected variety over an algebraically closed field k with the property that $\pi_1^{\mathrm{ab}}(U) = 1$ and such that it admits a good compactification X, where $X - U = \cup_{i\in I} Z_i =: Z$ is a simple strict normal crossings divisor. Denote by W the Witt ring of k and by K the fraction field of W.

We endow X with the log structure associated to the strict normal crossings divisor Z and denote this log scheme by (X, M_Z) or (X, Z): if we denote the open immersion $X \setminus Z \hookrightarrow X$ by j, the inclusion

$$M_Z := \mathcal{O}_X \cap j_* \mathcal{O}_{X-Z}^\times \hookrightarrow \mathcal{O}_X \tag{1.1}$$

defines a log structure on X.

Definition 1.1 (Trivial log structure)**.** *[Sh00, Example 2.4.1].*

Let Y be a scheme or a p-adic formal scheme. The pair $(\mathcal{O}_Y^\times, \mathcal{O}_Y^\times \hookrightarrow \mathcal{O}_Y)$ is a log structure on Y. This log structure is associated to the pre-log structure $\{1\}_Y \to \mathcal{O}_Y$.

Since Z is a normal crossings divisor, we know by [Sh00, Example 2.4.4] that M_Z is an fs, hence also fine, log structure [Sh00, Definition 2.1.5] and the morphism $(X, M) \to (\mathrm{Spec}\, k, \text{triv. log. str.})$ is log smooth.

Definition 1.2. *[Sh02],[HyKa] Assume that we have a morphism of fine log schemes*

$$(X, M) \to (Spf\, V, N)$$

with $X/\operatorname{Spf} V$ of finite type and V a totally ramified finite extension of W.

A good embedding system is a diagram

$$\begin{array}{ccc} (X^\bullet, M^\bullet) & \longrightarrow & (P^\bullet, N^\bullet) \\ \downarrow & & \downarrow \\ (X, M) & \longrightarrow & (Spf\, V, N) \end{array} \tag{1.2}$$

with the following properties:

1. *$X^i \to X$ is a hyper-covering for the étale topology and $M^i (i \geq 0)$ is the inverse image of M on X^i, for all i.*
2. *each (X^i, M^i) is a simplicial fine log scheme, of Zariski type and of finite type over k.*
3. *each (P^i, N^i) is a simplicial fine formal log V scheme, formally log smooth and of Zariski type.*
4. *$(X^i, M^i) \to (P^i, N^i)$ is a closed immersion for all i.*

Remark 1.3. *By [Sh02, Propostion 2.2.11], given morphisms of fine log schemes $(X, M) \xrightarrow{f} (\mathrm{Spec}\, k) \xrightarrow{i} (Spf\, V, N)$ where f is of finite type, i is the canonical exact closed immersion and assuming that $(Spf\, V, N)$ admits a chart, there exists at least one good embedding system of (X, M) over $(Spf\, V, N)$.*

In [NaSh08, §2.2] they define a log version of classical crystalline linearization functor: let S be a fine log scheme with a fine PD structure on a quasicoherent ideal sheaf $\mathcal{I}$ of $\mathcal{O}_S$ with the property that a power of p kills $\mathcal{O}_S$ (e.g. $S = W_n$). Let $Y \to S$ be a morphism of fine log schemes such that the PD structure on S extends to (the underlying scheme of) Y.

For an object T of the log crystalline site $(Y/S)^{log}_{crys}$ we have the localization functor

$$j_T : \widetilde{(Y/S)^{log}_{crys}}|_T \to (Y/S)^{log}_{crys}$$

and the morphism of ringed topoi

$$\phi : (\widetilde{(Y/S)^{log}_{crys}}|_T, \mathcal{O}_{Y/S}|_T) \to (\widetilde{T}_{\mathrm{Zar}}, \mathcal{O}_T) \tag{1.3}$$

which is defined as follows: for an $\mathcal{O}_T$-module $\mathcal{E}$ and an object $T' \xrightarrow{\psi} T$ in the localization, we set $\phi^*(\mathcal{E})(T', \psi) := \Gamma(T', \psi^*(\mathcal{E}))$ (denote by T, T' the sheaf on the crystalline site representable by T, T' respectively).

Let $i : Y \hookrightarrow \mathcal{Y}$ be an embedding into a smooth log scheme over S to which the PD structure of S extends. We can take the element $(Y, \mathcal{D}_Y(\mathcal{Y}))$ of $(Y/S)^{log}_{crys}$, where

$\mathcal{D}_Y(\mathcal{Y})$ is the log PD envelope of i. We then have a commutative diagram of topoi [NaSh08, (2.2.0.4)]

$$\begin{array}{ccc} \widetilde{(Y/S)^{log}_{crys}}|_T & \xrightarrow{j_T} & \widetilde{(Y/S)^{log}_{crys}} \\ \downarrow \phi & \swarrow_{u_{Y/S}} & \\ \widetilde{\mathcal{D}_Y(\mathcal{Y})}_{\mathrm{Zar}} = \widetilde{Y}_{\mathrm{Zar}} & & \end{array} .$$

Then the log linearization functor is defined as

$$L(\mathcal{E}) := j_{\mathcal{D}_Y(\mathcal{Y})*}\phi^*(\mathcal{E}) \in \widetilde{(Y/S)^{log}_{crys}}$$

and can be considered as the exact functor [NaSh08, Lemma 2.2.4]

$$\{\text{category of } \mathcal{O}_{\mathcal{D}_Y(\mathcal{Y})} - \text{modules and } \mathcal{O}_{\mathcal{D}_Y(\mathcal{Y})} - \text{linear morphisms}\} \to$$

$$\to \{\text{the category of crystals of } \mathcal{O}_{Y/S} - \text{crystals}\}.$$

Proposition 1.4. *[NaSh08, Proposition 2.2.7]*
With the above notation, the natural morphism

$$\mathcal{O}_{Y/S} \to L(\mathcal{O}_{\mathcal{D}_Y(\mathcal{Y})} \otimes_{\mathcal{O}_{\mathcal{Y}}} \omega^{\bullet,log}_{\mathcal{Y}/S}) \tag{1.4}$$

is a quasi isomorphism.

In the rest we freely use the terminology and notation of [ES16] and [Sh02].

2 Chern classes

Recall that by [ES16, Proposition 2.10] a rank 1 isocrystal admits a locally free lattice. We can now use the previous constructions to prove the following:

Lemma 2.1. *Let L be a locally free lattice of a rank one isocrystal $\mathcal{L}$ on U, which extends to a log isocrystal on X. Then $c_1^{logcrys}(L_X) = 0$.*

Proof. Let

$$\begin{array}{ccc} (X^\bullet, M^\bullet) & \longrightarrow & (P^\bullet, N^\bullet) \\ \downarrow & & \downarrow \\ (X, M_Z) & \longrightarrow & (\mathrm{Spf}\ W_n, N) \end{array} \tag{2.1}$$

be a good embedding for the scheme $X = \overline{U}$ equipped with the log structure associated to the sncd Z, over Spf W_n where N is the trivial log structure (and therefore

admits a chart). Denote by $D_n^\bullet$ the completed PD envelope of $X^\bullet \hookrightarrow P^\bullet$ and by i the map $i : X^\bullet \to D_n^\bullet$. We can apply the previous discussion to $X^\bullet \hookrightarrow P^\bullet$ over $S = \operatorname{Spf} W_n$ and restricting (1.4) to the crystalline units we get

$$Ru_{X^\bullet/W_n *}\mathcal{O}^\times_{X^\bullet/W_n} \simeq (\mathcal{O}^\times_{D_n^\bullet} \xrightarrow{dlog} \omega^{1,log}_{D_n^\bullet} \to \omega^{2,log}_{D_n^\bullet} \to \cdots) \tag{2.2}$$

in $\widetilde{(X)}_{\mathrm{Zar}}$.

One defines the log crystalline Chern classes, as remarked also in [ABV05, §6], in complete analogy to the crystalline Chern classes of [BI70]. Adapted to the present situation, the construction of these classes is as follows: define $\mathcal{J}_{X^\bullet/W_n}$ by the exact sequence in $(\widetilde{X^\bullet/W_n})$

$$0 \to \mathcal{J}_{X^\bullet/W_n} \to \mathcal{O}_{X^\bullet/W_n} \to i_*\mathcal{O}_{X^\bullet} \tag{2.3}$$

which in turn yields

$$1 \to 1 + \mathcal{J}_{X^\bullet/W_n} \to \mathcal{O}^\times_{X^\bullet/W_n} \to i_*\mathcal{O}^\times_{X^\bullet}. \tag{2.4}$$

Then $\mathcal{J}_{X^\bullet/W_n}$ is a log PD ideal, so one defines the logarithm $1 + \mathcal{J}_{X^\bullet/W_n} \to \mathcal{J}_{X^\bullet/W_n}$, $x \mapsto log(1+x)$.

Then the log crystalline Chern class $c_1^{logcrys} : H^1(X, \mathcal{O}_X^\times) \to H^2(X/W_n)$ is computed by

$$\begin{aligned} &H^1(X, \mathcal{O}_X^\times) \to H^1(X^\bullet, \mathcal{O}^\times_{X^\bullet}) \to H^2((X^\bullet/W_n)^{log}_{crys}, 1 + \mathcal{J}_{X^\bullet/W_n}) \xrightarrow{log} \\ &\to H^2((X^\bullet/W_n)^{log}_{crys}, \mathcal{J}_{X^\bullet/W_n}) \to H^2(X^\bullet/W_n)^{log}_{crys} = H^2(X/W)^{log}_{crys}. \end{aligned}$$

On the other hand, we can compute the Chern classes also via the completed PD envelope. Applying $Ru_{X^\bullet/W_n *}$ to (2.4) and using (2.2) we get an exact sequence in $\widetilde{(D_n^\bullet)}_{Zar}$

$$1 \to ((1 + J^\bullet) \to \omega^{\geq 1,log}_{D_n^\bullet/W_n}) \to (\mathcal{O}^\times_{D_n^\bullet} \xrightarrow{dlog} \omega^{\geq 1,log}_{D_n^\bullet/W_n}) \to \mathcal{O}_X^\times \to 1 \tag{2.5}$$

where $J^\bullet = \operatorname{Ker}(\mathcal{O}^\times_{D^\bullet} \to \mathcal{O}_X^\times)$. We can then compute the log crystalline Chern class by

$$\begin{aligned} &H^1(X, \mathcal{O}_X^\times) \to H^1(X^\bullet, \mathcal{O}^\times_{X^\bullet}) \to H^2((D_n^\bullet/W_n)^{log}_{crys}, (1 + J^\bullet) \to \omega^{\geq 1,log}_{D_n^\bullet/W_n}) \xrightarrow{log} \\ &\to H^2((D_n^\bullet/W_n)^{log}_{crys}, J^\bullet \xrightarrow{d} \omega^{\geq 1,log}_{D_n^\bullet/W_n}) \to H^2(D_n^\bullet/W_n)^{log}_{crys} = H^2(X/W_n)^{log}_{crys}, \end{aligned}$$

with the last equality being true by [HyKa, Proposition 2.10] .

Since L_X is the value on X of a locally free log crystal on X, this is equivalent to an $\mathcal{O}_D$-module with quasi nilpotent integrable log connection and its Chern class is zero in $H^2(X/W_n)^{log}_{crys}$ for all n, hence also when the Chern class map is seen as $c_1 : H^1(X, \mathcal{O}_X^\times) \to H^2(X/W)^{log}_{crys}$, the Chern class of L_X is zero. □

Lemma 2.2. *The line bundle $L_X^{\otimes N}$ for some natural number N can be written as $\mathcal{O}_X(\sum a_i Z_i)$, with $a_i \in \mathbb{Q}$.*

Proof. We have an exact sequence:

$$0 \to H^0(X, \mathcal{O}_X^\times) \to H^0(U, \mathcal{O}_U^\times) \to \oplus \mathbb{Z} Z_i \to \mathrm{Pic}(X) \to \mathrm{Pic}(U) \to 0 \tag{2.6}$$

By [Kin13, Proposition 3.5] we have that $\pi_1^{\mathrm{ab}}(X) = \pi_1^{\mathrm{ab}}(U) = 1$ implies $H^0(U, \mathcal{O}_U^\times) = k^\times$ and thus $H^0(X, \mathcal{O}_X^\times)$ is isomorphic to $H^0(U, \mathcal{O}_U^\times)$.

Moreover, it also implies that $\mathrm{Pic}(X) = \mathrm{NS}(X)$ and $\mathrm{Pic}(U) = \mathrm{NS}(U)$, and the exact sequence becomes

$$0 \to \oplus \mathbb{Z} Z_i \to \mathrm{NS}(X) \to \mathrm{NS}(U) \to 0. \tag{2.7}$$

The log crystalline Chern class can be seen as

$$c_1^{logcrys} : H^1(X, \mathcal{O}_X^\times) \to H^2(X/W)^{log}_{crys} \otimes \mathbb{Q} \simeq H^2_{rig}(U/K). \tag{2.8}$$

We observe that the following diagram commutes:

$$\begin{array}{ccc} H^1(X, \mathcal{O}_X^\times) & \xrightarrow{c_1^{logcrys}} & H^2(X/W)^{log}_{crys} \\ \downarrow{\scriptstyle c_1^{crys}} & \nearrow{\scriptstyle f^*} & \\ H^2(X/W)_{crys} & & \end{array} , \tag{2.9}$$

where we denote by $f^* : H^2(X/W)_{crys} \to H^2(X/W)^{log}_{crys}$ the homomorphism which is obtained by the exact map of topoi $f : (X/W_n)^{log}_{crys} \to (X/W_n)_{crys}$ for all n and is defined as $f^*(E)((U, M_U), (T, M_T)) := E(U \subset T)$. As stated in [ABV05] this map is exact and commutes with global sections. Applying this to the sequence

$$1 \to 1 + \mathcal{J}_X \to \mathcal{O}^\times_{X/W_n} \to \mathcal{O}_X^\times \to 1 \tag{2.10}$$

we get

$$\begin{array}{ccccccccc} 1 & \longrightarrow & f^*(1+\mathcal{J}_X) & \longrightarrow & f^*(\mathcal{O}^\times_{X/W_n}) & \longrightarrow & f^*(\mathcal{O}_X^\times) & \longrightarrow & 1 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 1 & \longrightarrow & 1+\mathcal{J}_X^{log} & \longrightarrow & \mathcal{O}^{\times,log}_{X/W_n} & \longrightarrow & \mathcal{O}_X^\times & \longrightarrow & 1 \end{array} \tag{2.11}$$

with exact rows and commutative squares. By taking cohomology of the first row and composing with the log map, one obtains $f^*(c_1^{crys})$ whereas by taking cohomology of the second, one obtains $c_1^{logcrys}$.

Combining the previous observation with [Pet01, Proposition 3.6.1] we obtain the following commutative diagrams

$$\begin{array}{ccc} H^2_{rig}(X/K) & \longrightarrow & H^2(X/W)_{crys} \otimes \mathbb{Q} \\ {\scriptstyle c_1^{rig}}\uparrow & \overset{c_1^{crys}}{\nearrow} & \downarrow \\ H^1(X, \mathcal{O}^\times) & \xrightarrow[c_1^{logcrys}]{} & H^2(X/W)^{log}_{crys} \otimes \mathbb{Q} \end{array} . \tag{2.12}$$

Moreover, by [Ill79, Remark II 6.8.4], and using the fact that $\pi_1^{\mathrm{ab}}(X) = 1$ implies also that $\mathrm{Pic}(X) = \mathrm{NS}(X)$, we know that $H^1(X, \mathcal{O}_X^\times) \otimes K \hookrightarrow H^2_{rig}(X/K)$ via the first rigid Chern class.

Our line bundle L_X seen in $\mathrm{NS}(X) \otimes \mathbb{Q}$ (i.e. L_X modulo torsion) is sent to zero by $c_1^{logcrys}$, by Lemma 2.1. So we have

$$\begin{array}{ccccc} & & & & \langle KZ_i \rangle_{\mathbb{Q}} \\ & & & & \downarrow \\ \mathrm{NS}(X) \otimes \mathbb{Q} & \hookrightarrow & \mathrm{NS}(X) \otimes K & \xrightarrow{c_1^{rig}} & H^2_{rig}(X/K) \\ & & & \overset{j^* c_1^{rig}}{\searrow} & \downarrow j \\ & & & & H^2_{rig}(U/K) \end{array}$$

which means thatL_X is contained in the kernel of the map $H^2_{rig}(X/K) \to H^2_{rig}(U/K)$, which is $\langle KZ_i \rangle_{\mathbb{Q}}$. Hence, $L_X \in \langle KZ_i \rangle_{\mathbb{Q}} \cap \mathrm{NS}(X) \otimes \mathbb{Q} = \oplus \mathbb{Q} Z_i$.

This in turn means that L_X can be written as $\mathcal{O}_X(\sum a_i Z_i) \otimes T$, with T a torsion bundle, and therefore some power of it can be written as a formal sum of Z_i with $\mathbb{Q}$-coefficients. □

3 Residue exact sequence

We follow [ABV05, §6]. Let $\{U_i \hookrightarrow V_i\}$ be a covering in the log crystalline site. We set $U_{ij} = U_i \cap U_j$ and denote by V_{ij} the log PD envelope of U_{ij} in V_i (or V_j) and do the same for every tuple of indices $(i_0, i_1, \cdots, i_n)$. Define $Q^r_{i_0,\ldots,i_n} := \omega^{r,log}_{V_{i_0,\ldots,i_n}} / \omega^r_{V_{i_0,\ldots,i_n}}$. We then have the following maps between the total complexes of the corresponding Čech complexes, with exact columns:

(3.1)

$$\begin{array}{ccccccccc}
0 & \longrightarrow & \oplus\mathcal{O}^{\times}_{V_i} & \longrightarrow & \oplus_{i,j}\mathcal{O}^{\times}_{V_{ij}} \oplus_i \omega^1_{V_i} & \longrightarrow & \oplus_{i,j,k}\mathcal{O}^{\times}_{V_{ijk}} \oplus_{i,j} \omega^1_{V_{ij}} \oplus_{i,j} \omega^2_{V_{ij}} & \longrightarrow & \dots \\
 & & \downarrow & & \downarrow & & \downarrow & & \\
0 & \longrightarrow & \oplus\mathcal{O}^{\times}_{V_i} & \longrightarrow & \oplus_{i,j}\mathcal{O}^{\times}_{V_{ij}} \oplus_i \omega^{1,log}_{V_i} & \longrightarrow & \oplus_{i,j,k}\mathcal{O}^{\times}_{V_{ijk}} \oplus_{i,j} \omega^{1,log}_{V_{ij}} \oplus_{i,j} \omega^{2,log}_{V_{ij}} & \longrightarrow & \dots \\
 & & \downarrow & & \downarrow & & \downarrow & & \\
 & & 0 & \longrightarrow & \oplus Q^1_i & \longrightarrow & \oplus_{i,j} Q^1_{ij} \oplus_i Q^2_i & \longrightarrow & \dots \\
 & & & & \downarrow & & \downarrow & & \\
 & & & & 0 & & 0 & &
\end{array}$$

and note that the third complex is the total complex of:

(3.2)

$$\begin{array}{ccccccc}
0 & \longrightarrow & \oplus Q^1_i & \longrightarrow & \oplus Q^1_{ij} & \longrightarrow & \dots \\
\downarrow & & \downarrow & & \downarrow & & \\
0 & \longrightarrow & \oplus Q^2_i & \longrightarrow & \oplus Q^2_{ij} & \longrightarrow & \dots \\
\downarrow & & \downarrow & & \downarrow & & \\
0 & \longrightarrow & \oplus Q^3_i & \longrightarrow & \oplus Q^3_{ij} & \longrightarrow & \dots \\
\downarrow & & \downarrow & & \downarrow & & \\
0 & & \dots & & \dots & & \dots
\end{array}$$

Taking cohomology, we obtain maps (starting from H^1)

(3.3)
$$H^1_{crys}(X/W, \mathcal{O}^{\times}_{X/W}) \to H^1((X,M)/W, \mathcal{O}^{\times}_{X/W})^{log}_{crys} \to H^1(Tot(Q_{\bullet,\bullet})) \to H^2(X/W, \mathcal{O}^{\times}_{X/W})$$

and computing the cohomology of the complex (3.2) with spectral sequences, using the fact that $H^1(Tot(Q_{\bullet,\bullet})) = horE^{\infty}_{10} \oplus horE^{\infty}_{01}$, we obtain the exact sequence

$$\begin{aligned}(3.4)\quad H^1_{crys}(X/W, \mathcal{O}^{\times}_{X/W}) \to H^1((X,M)/W, \mathcal{O}^{\times}_{X/W})^{log}_{crys} \to \\ \to \text{Ker } (H^0(Q^1_{\bullet}) \to H^0(Q^2_{\bullet})) \to H^2(X/W, \mathcal{O}^{\times}_{X/W}).\end{aligned}$$

By the computation in [ABV05, Proposition 6.4] we have that

$$(3.5)\qquad \text{Ker } (H^0(Q^1_{\bullet}) \to H^0(Q^2_{\bullet})) = \text{Div}_Z(X) \otimes W(k), \text{ or } \oplus W(k)[Z_i],$$

so we obtain:

(3.6)
$$H^1_{crys}(X/W, \mathcal{O}^{\times}_{X/W}) \to H^1((X,M)/W, \mathcal{O}^{\times}_{X/W})^{log}_{crys} \to \oplus W(k)[Z_i] \to H^2(X/W, \mathcal{O}^{\times}_{X/W}).$$

Moreover, in [ABV05] we find the exact sequence

$$0 \to H^1_{crys}(X/W) \to H^1((X,M)/W)^{log}_{crys} \to \oplus W(k)[Z_i] \to H^2_{crys}(X/W) \tag{3.7}$$

with the last map being the crystalline Chern class map. By [ES16, Proposition 2.9(2)] and [Esn16, Theorem 4.3.1] we have that $\pi_1^{\mathrm{ab}}(X) = 1$ implies $H^1_{crys}(X/W) = 0$. Hence, the group $H^1(X/W)^{log}_{crys}$ coincides in this case with the kernel of the crystalline Chern class map, which by [ABV05, Proposition 6.4] is equal to $\mathrm{Div}^0_Z(X) \otimes W(k)$ and this is equal to zero because $\mathrm{Pic}^0(X) = 0$. We therefore obtain an injection $\oplus W(k)[Z_i] \hookrightarrow H^2_{crys}(X/W)$.

For X proper we have that $H^1_{crys}(X/W, \mathcal{O}_X^\times) = 0$ by [ES16, Theorem 0.1.1] and [Esn16, Theorem 4.3.1], and (3.6) then gives us an injection

$$H^1((X,M)/W, \mathcal{O}^\times)^{log}_{crys} \hookrightarrow \oplus W(k)[Z_i]. \tag{3.8}$$

Combining the two exact sequences we obtain an injection:

$$H^1((X,M)/W, \mathcal{O}^\times)^{log}_{crys} \hookrightarrow \oplus W(k)[Z_i] \hookrightarrow H^2_{crys}(X/W). \tag{3.9}$$

Furthermore, by Lemma 2.2 one has that under our assumptions the crystal $E := L^{\otimes N}$ for some N lies in $\oplus \mathbb{Q}[Z_i]$.

From the exact sequences above and going through the definition of c_1^{crys} and $c_1^{logcrys}$ we observe, as already remarked in (Ibid.), that the following diagram commutes:

$$\begin{array}{ccccc} H^1((X,M)/W, \mathcal{O}^\times)^{log}_{crys} & \hookrightarrow & \oplus W(k)[Z_i] & \xrightarrow{c_1^{crys}} & H^2(X/W)_{crys} \\ & \searrow & & \nearrow_{c_1^{crys}} & \\ & & H^1(X, \mathcal{O}_X^\times) & & \end{array} . \tag{3.10}$$

From this we conclude that under our assumptions the projection map

$$H^1((X,M)/W, \mathcal{O}^\times)^{log}_{crys} \to H^1(X, \mathcal{O}_X^\times), \qquad E \mapsto E_X \tag{3.11}$$

is in fact injective, and so the value of E on X can also be written as a sum with $W(k)$ coefficients.

Hence, E_X lies in $\oplus \mathbb{Z}_{(p)}[Z_i]$. Therefore there is $m \in \mathbb{Z} \setminus p\mathbb{Z}$ with the property that the bundle $E_X^{\otimes m} \in \oplus \mathbb{Z}[Z_i]$ is such that $E_U^{\otimes m} = \mathcal{O}_U$ and $(m,p) = 1$. This in turn defines a Kummer cover of U, for which by assumption we have that $\pi_1^{\mathrm{tame,ab}}(U) = 1$. Hence m can be chosen as 1 and E_X is in fact in $\oplus \mathbb{Z}[Z_i]$ and has the property that E_U is trivial (see Lemma 4.2 for detailed argument).

Remark 3.1. *We can show that $H^1_{rig}(U/K) = 0$ under our assumptions. For a proper scheme this is known. Observing (3.7) we conclude as before that $H^1(X/W)^{log}_{crys} = 0$. Hence $H^1_{rig}(U/K) \simeq H^1(X/W)^{log}_{crys} \otimes \mathbb{Q} = 0$.*

Lemma 3.2. *Suppose $E_U \simeq \mathcal{O}_U$. Then we can choose a locally free extension $E_X \simeq \mathcal{O}_X$.*

Proof. If $E_U \simeq \mathcal{O}_U$, then $E_X \simeq \mathcal{O}_X(\sum b_i Z_i)$, with $E_X \xrightarrow{\nabla} E_X \otimes \Omega^1(\log Z)$. Let j be the map $U \to X$. Inside $\mathcal{O}_X = j_*\mathcal{O}_U \xrightarrow{d} j_*\Omega^1_U = \Omega^1_X(\log Z)$ we have $\mathcal{O}_X(-a_iZ_i) \to \mathcal{O}_X(-a_iZ_i) \otimes \Omega^1(\log Z)$, since $d(t_i^{a_i}.1) = a_i \frac{t_i^{a_i}}{t_i}.\frac{dt_i}{t_i} \in \mathcal{O}_X(-a_iZ_i) \otimes \Omega^1(\log Z)$.

We can write the connection ∇ as $d + \alpha, \alpha \in H^0(X, \Omega^1(\log Z))$. Then writing E_X as $\mathcal{O}_X(\sum a_iZ_i - \sum n_iZ_i)$ with $a_i, n_i > 0$, we have:

$$\begin{array}{ccc} \mathcal{O}_X(\sum a_iZ_i - \sum n_iZ_i) & \longrightarrow & \mathcal{O}_X(\sum n_iZ_i) \\ \uparrow & & \uparrow \\ \mathcal{O}_X(-\sum n_iZ_i) & \longrightarrow & \mathcal{O}_X \end{array} \tag{3.12}$$

and the connections:

$$\begin{array}{ccc} E_X & \longrightarrow & E_X \otimes \Omega^1(\log Z) \\ \uparrow & & \\ \mathcal{O}_X(-\sum n_iZ_i) & \longrightarrow & \mathcal{O}_X(-\sum n_iZ_i) \otimes \Omega^1(\log Z) \\ \downarrow & & \\ \mathcal{O}_X & \longrightarrow & \mathcal{O}_X \otimes \Omega^1(\log Z) \end{array} \tag{3.13}$$

So, for E crystal on U/k such that $E_U = \mathcal{O}_U$ we can choose $E_X \simeq \mathcal{O}_X$. □

4 Triviality of isocrystal

Theorem 4.1. *Let E a locally free lattice of a rank 1 isocrystal $\mathcal{E}$ on U that extends to a log crystal on X and such that E_X is trivial. Then E is trivial.*

Proof. **Step 1**: We have the following maps of fine log schemes:

$$(X, Z) \to \underline{S} \to \underline{W}(k)$$

where $\underline{S}$ is the log scheme defined by $S = \operatorname{Spec} k$ with the log structure associated to $\mathbb{N} \to k; 1 \mapsto 0$ and $\underline{W}(k)$ is the log scheme defined by $\operatorname{Spec} W(k)$ with the log structure $\mathbb{N} \to W(k); 1 \mapsto 0$.

There is at least one good embedding system for $(X, Z)/(\operatorname{Spec} k, N)$, so we assume from now on that there is an embedding $(X, Z) \hookrightarrow (P, L)$ to a smooth formal scheme over $(\operatorname{Spf} W, N)$.

Denote by D the completed log PD envelope of $(X \hookrightarrow P)$. It is the usual PD envelope equipped with the inverse image of the log structure. Denote by D_n the

corresponding PD envelopes for the immersions over $P/\operatorname{Spf} W_n$.

$$\begin{array}{ccc} D_n & & \\ \uparrow & \searrow & \\ X & \longrightarrow & P_{W_n} \\ \downarrow & \swarrow & \\ W_n & & \end{array} \cdot \tag{4.1}$$

By [Ka89, Theorem 6.2] the category $C_{crys}(X/W_n)^{log}$ of log crystals on X/W_n is equivalent to the category of $\mathcal{O}_{\mathcal{D}_n}$-modules with integrable log quasi nilpotent connection.

Let E be a log crystal such that the restriction E_n of E_{n+1} to $C_{crys}(X/W_n)^{log}$ is the trivial log crystal. We can define, as in [ES16, Proposition 3.6], $\mathcal{D}$ to be the set of pairs (G, ϕ) such that $G \in C_{crys}(X/W_{n+1})^{log}$ with $\phi : \mathcal{O}_X \simeq \overline{G} \in C_{crys}(X/W_n)^{log}$. We claim that in this case we have an isomorphism $e : \mathcal{D} \simeq H^1(X/W_1)^{log}_{crys}$. By the equivalence of log crystals and modules with log integrable quasi nilpotent connection, we can identify E_{n+1} with a $\mathcal{O}_{\mathcal{D}_{n+1}}$ module M_{n+1} with a log connection ∇_{n+1}. By assumption, there is an isomorphism

$$\phi : (\mathcal{O}_{D_n}, d) \to (M_n, \nabla_n),$$

(M_n, ∇_n) being the restriction to $MIC^{log}(D_n)^{qn} \simeq C_{crys}(X/W_n)^{log}$.

Take an open affine cover $U = \{U_a\}$ of D_{n+1} and an isomorphism $\psi_a : \mathcal{O}_{U_a} \to M_{n+1}|_{U_a}$ lifting $\phi|_{D_n \times U_a}$. Then $\psi_a^*(\nabla_{n+1})$ defines a connection on $\mathcal{O}_{Ua}$ that is given as $d + p^n s_a$ with $s_a \in \Gamma(U_a, \omega^1_{log,D_1})$. On the intersection $U_a \cap U_b =: U_{ab}$ the glueing $(\psi_a|_{U_{ab}})^{-1} \circ (\psi_b|_{U_{ab}})$ is given by $1 + p^n z_{ab}$ with $z_{ab} \in \Gamma(U_{ab}, \mathcal{O}_{D_1})$.

Because of the integrability of the connection and the compatibility of the connection with the glueing, we see as in [ES16], that $(\{s_a\}, \{z_{ab}\})$ actually defines a 1-cocycle in Tot $\Gamma(U, \omega^\bullet_{log,D_1})$ and this defines a class in the cohomology $H^1(\text{Tot } \Gamma(U, \omega^\bullet_{log,D_1})) = H^1(X/W_1)^{log}_{crys}$.

Step 2: By the previous step we get:

$$Ker(MIC^{log}(D_{n+1}) \to MIC^{log}(D_n)) \simeq H^1(X, \omega^\bullet_{log,D_1}).$$

We know the following from [Sh02, Theorem 2.4.4 and Corollary 2.3.9], denoting by $K_{X/W}$ the convergent isocrystal defined by $T \mapsto K \otimes_W \Gamma(T, \mathcal{O}_T)$: for all $i \in \mathbb{N}$

$$H^i((X/W)^{log}_{crys}, K \otimes \mathcal{O}_{X/W}) \simeq H^i((X/W)^{log}_{conv}, K_{X/W}) \tag{4.2}$$

and

$$H^i((X/W)^{log}_{conv}, K_{X/W}) \simeq H^i_{rig}(U/K). \tag{4.3}$$

In [Sh02, page 136] Shiho defines for $\mathcal{E} = K \otimes \mathcal{F}$ in $\mathcal{I}_{crys}(X/W)^{log}$:

$$H^1((X/W)^{log}_{crys}, \mathcal{E}) = \mathbb{Q} \otimes_{\mathbb{Z}} H^1((X/W)^{log}_{crys}, \mathcal{F}).$$

Combining this with Remark 3.1

$$H^1((X/W)^{log}_{crys}, K \otimes \mathcal{O}_{X/W}) = \mathbb{Q} \otimes H^1(X/W)^{log}_{crys}, \mathcal{O}_{X/W}) = H^1_{rig}(U/K) = 0.$$

From [ABV05, Theorem B' and Section 2.3], we know that $H^1(X/W)^{log}_{crys}$ is a free W-module, hence torsion free and therefore $H^1(X/W)^{log}_{crys} = \varprojlim_n H^1(X/W_n)^{log}_{crys} = 0$. Hence there is a natural number N large enough such that for all $n \geq N$, the maps $H^1(X/W_n)^{log}_{crys} \to H^1(X/W_1)^{log}_{crys}$ are zero.

Claim: : There is a natural number $d \geq 0$ s.t. $(F^d)^* H^1(X/W_1)^{log}_{crys} = 0$.

From [Mum70, Corollary on page 143] we have the decomposition of $H^1(X/W_1)^{log}_{crys}$ as a direct sum of the part on which Frobenius acts as an isomorphism and the part on which it acts nilpotently. Denote the former by $H^1((X/W_1)^{log}_{crys})_{ss}$. We have the following maps:

$$\begin{array}{ccccc} H^0(X, \Omega_X^{\geq 1}) & \hookrightarrow & H^1_{crys}(X/k) & \longrightarrow & H^1(X, \mathcal{O}_X) \\ \downarrow & & \downarrow & & \downarrow \\ H^0(X, \omega^{\geq 1}_{log,X}) & \hookrightarrow & H^1(X/W_1)^{log}_{crys} & \longrightarrow & H^1(X, \mathcal{O}_X) \end{array} .$$

Since F^* acts by zero on the image of $H^0(X, \Omega^1_X)$ in $H^1_{crys}(X/k)$, it acts by zero on the image of $H^0(X, \omega^1_{log,X})$ in $H^1(X/W_1)^{log}_{crys}$ and therefore $H^1((X/W_1)^{log}_{crys})_{ss} \subset H^1(X, \mathcal{O}_X)_{ss} = 0$, the latter equality being true because:

$$H^1(X, \mathcal{O}_X)_{ss} = H^1(X, \mathcal{O}_X)^{F=1} \otimes_{\mathbb{F}_p} k = H^1_{\text{ét}}(X, \mathbb{F}_p) \otimes k = \text{Hom}(\pi_1^{\text{ab}}(X), \mathbb{F}_p) \otimes k = 0.$$

So, there exists some $d \in \mathbb{N}$ s.t. $(F^d)^* H^1(X/W_1)^{log}_{crys} = 0$.

By the same computations as in Step 1, one can actually have:

$$\text{Ker}\left(MIC^{log}(D_{n+m}) \to MIC^{log}(D_n)\right) \simeq H^1(X, \omega^\bullet_{log,D_m}), \text{ for all } 1 \leq m \leq n.$$

Assume now that we have E as in the assumption of the theorem. Then $E_2 \in \text{Ker}\left(MIC^{log}(D_2) \to MIC^{log}(D_1)\right)$, and this defines a class $e(E_2) \in H^1(X/W_1)^{log}_{crys}$. By the definition of d above, we have $(F^d)^* e(E_2) = e((F^*)^d E)_2 = 0$. Hence, by Step 1, $((F^*)^d E)_2$ is trivial. Repeating this we find $r \in \mathbb{N}$ such that $((F^r)^* E)_N = (F^r)^* E_N =: E'_N$ is trivial.

We also see that the image of E'_{2N} via the restriction $H^1(X, \omega^\bullet_{log,D_N}) \to H^1(X, \omega^\bullet_{log,D_1})$, which is equal to E'_{N+1} is trivial.

Continuing like this, we can show that E'_n is trivial for all $n \geq N$, therefore E' is trivial. The functor F^* is fully faithful restricted to locally free crystals, so we conclude that E is itself trivial.

□

This proves, setting as before $E = L^{\otimes N}$, using Lemma 3.2 and the conclusion of the discussion of the previous Section, that a power of the rank one crystal L is trivial. Using the following two lemmas, we conclude that L itself is trivial and the rank 1 isocrystal we started with is also trivial.

Lemma 4.2. *With the previous assumptions, if $L^{\otimes n} = 1$ with $(n, p) = 1$, then L is trivial.*

Proof. Since $L^{\otimes n}$ is trivial, we have $L_U^{\otimes n} \simeq \mathcal{O}_U$ with $(n, p) = 1$, hence

$$\underline{\mathrm{Spec}} \left(\bigoplus_{i=0}^{n-1} L_U^i \right) \to U$$

is a Kummer cover.

Since $\pi_1^{\mathrm{ab,tame}}(U) = 1$ this cover has to be trivial, so $n = 1$ and $L_U \simeq \mathcal{O}_U$. Then we can choose a locally free extension L_X that is trivial. Applying the previous theorem to this, we get the desired result. □

Again with the previous notation we have the following:

Lemma 4.3. *If $L^{\otimes p} = 1$, L is trivial.*

Proof. If $L^{\otimes p} = 1$ then as coherent sheaves $L_U^{\otimes p} \simeq \mathcal{O}_U$, thus $M_X := (F^*L)_U \simeq \mathcal{O}_U$. We choose a locally free extension M_X that is the trivial log crystal and apply Theorem 4.1 (F^*L is also log extendable if L is). We get that F^*L is trivial. But the functor $F^* : \mathrm{Crys}(U/W) \to \mathrm{Crys}(U/W)$ restricted to locally free crystals is fully faithful [Ogu94, Ex. 7.3.4]. Hence L itself is trivial. □

Corollary 4.4. *Extensions of the trivial object by itself in $I_{crys}(X/W)^{log}$ are trivial. In particular, any isocrystal on U whose extension to X is of this form, is trivial.*

Proof. The group of log isocrystals $\mathcal{E}$ on $(X, M)/W$ for which

$$0 \to \mathcal{O}_{X/W} \otimes \mathbb{Q} \to \mathcal{E} \to \mathcal{O}_{X/W} \otimes \mathbb{Q} \to 0 \tag{4.4}$$

is exact, is precisely $\mathrm{Ext}^1(\mathcal{O}_{X/W} \otimes \mathbb{Q}, \mathcal{O}_{X/W} \otimes \mathbb{Q}) = H^1((X, M)/W, \mathcal{O}_{X/W} \otimes \mathbb{Q})^{log}_{crys}$. However,

$$H^1((X, M)/W, \mathcal{O}_{X/W} \otimes \mathbb{Q})^{log}_{crys} = H^1((X, M)/W, \mathcal{O}_{X/W})^{log}_{crys} \otimes \mathbb{Q} \simeq H^i_{rig}(U/K) = 0. \tag{4.5}$$

Let now $\mathcal{E}$ be an isocrystal on (U/W) that extends to a log isocrystal on X such that this extension fits into an exact sequence

$$0 \to \mathbb{Q}_{X/W} \to \mathcal{E} \to \mathbb{Q}_{X/W} \to 0 \tag{4.6}$$

in $I_{crys}(X/W)^{log}$. Then we also have an exact sequence in Crys $(U/W)_{\mathbb{Q}}$

$$0 \to \mathcal{O}_{U/W} \otimes \mathbb{Q} \to \mathcal{E} \to \mathcal{O}_{U/W} \otimes \mathbb{Q} \to 0. \tag{4.7}$$

Indeed, (4.6) being exact means that for all log PD thickenings $((V, M_V), (T, M_T))$ in the log crystalline site of X/W, we have

$$0 \to \mathcal{O}_T \otimes \mathbb{Q} \to \mathcal{E}_T \to \mathcal{O}_T \otimes \mathbb{Q} \to 0 \tag{4.8}$$

in Crys (U/W). The log PD thickenings $((V, M_V), (T, M_T))$ restrict to PD thickenings (V, T) on X and they in turn define PD thickenings $(V \cap U, T)$ on U. Every open of U is an open in X, thus we have the above exact sequence in Crys $(U/W)_{\mathbb{Q}}$. So $\mathcal{E}$ is itself an extension of the trivial isocrystal on U by itself and its extension to X is trivial. Applying the above theorem to this, we obtain that $\mathcal{E}$ is trivial on Crys $(U/W)_{\mathbb{Q}}$. □

5 Defining the residue map

We can in fact define the homomorphism $H^0((X, Z)/W, \mathcal{O}_X^\times) \to \oplus W(k)[Z_i]$ of the exact sequence (3.6) as a residue map assuming the existence of a lift. Assume we have as before, $U \hookrightarrow X = \overline{U} \hookleftarrow Z = \cup_i Z_i$ a sncd. Take an affine open X^0 of X and a smooth lifting X^0_W over W, and the corresponding lifting Z^0_W of $Z^0 = Z \cap X^0$. Denote the log PD envelope of $(X^0, Z^0) \hookrightarrow (X^0_W, Z^0_W)$ by D^0 (endowed with the pullback of the log structure on X^0).

Then a log crystal on X^0 is equivalent to the datum of an $\mathcal{O}_{D^0}$-module with an integrable log connection $\nabla : E_{D^0} \to E_{D^0} \otimes_{\mathcal{O}_{X^0_W}} \Omega^1_{X^0_W}(\log Z^0_W)$.

This log connection has well defined residues:

$$res_i : E_{D^0} \to \oplus \mathcal{O}_{(Z^0_W)^i} \otimes E_{D^0}$$

that are $\mathcal{O}_{X^0_W}$- linear maps. Because E has rank 1, these maps are actually elements of $H^0((Z^0_W)^i, \mathcal{O}_{(Z^0_W)^i})$.

Claim: For a smooth divisor $\Delta \subseteq Y$, for Y a smooth scheme over W, the residue map $E_Y \xrightarrow{res} \mathcal{O}_\Delta \otimes E_Y$, which is an element in $\Gamma(Y, \mathcal{O}_\Delta)$, is in particular a closed form.

Proof: We show that for $E_Y \xrightarrow{res} \mathcal{O}_\Delta \otimes E_Y \xrightarrow{d} \Omega^1_Y \otimes E_Y$, we have $d(res) = 0$: since E has rank 1, it can be written as $\mathcal{O}_Y e$ for some e such that $\nabla(e) = a\frac{dt}{t}$, $a \in \Gamma(Y, E_Y)$. Then $res(\nabla) = a_{|t=0}$.

Locally, choosing coordinates $(t, t_1, \cdots, t_r, p)$ such a has an expression as a sum $a = a_0 + a_1 t + \cdots$ with a_i functions on $(t_1, \cdots, t_r, p)$ and since the connection is integrable, we have that $0 = \nabla \circ \nabla(e) = \nabla(a\frac{dt}{t}) = da\frac{dt}{t} = (da_0 + t da_1 + \cdots)\frac{dt}{t}$. Restricting the last part of the equation to $t = 0$, we get $da_0 = da_{|t=0} = 0$. Hence $d(res) = da_{|t=0} = da_{0|t=0} = 0$ as claimed.

From [EV92, Chapter 8], we have for liftings over W_2: let X be a smooth scheme over $S := \text{Spec } k$, $Z = \sum_{j=1}^r Z_j$ ncd over S and $\tilde{Z}_j \subset \tilde{X}$ subschemes with $\tilde{Z}_j \otimes \tilde{S} = Z_j$, for $\tilde{S} = \text{Spec } W_2$. Then the components of $\tilde{Z}$ are Cartier divisors in $\tilde{X}$ if and only if $\tilde{Z} = \sum \tilde{Z}_j$ is a lifting of Z.

We can do the same steps to pass to a smooth lift from W_2 to W_3:

$$\begin{array}{ccccc} X & \longrightarrow & \tilde{X} & \longrightarrow & \tilde{X}' \\ \downarrow & & \downarrow & & \downarrow \\ \text{Spec } k & \longrightarrow & \text{Spec } W_2 & \longrightarrow & \text{Spec } W_3 \end{array} .$$

Then, if $\tilde{Z}' \subset \tilde{X}'$ a lifting of $\tilde{Z} \subset \tilde{X}$, the flatness of $\tilde{X}'$ and $\tilde{Z}'$ (by definition of the lifting) implies that the ideal $J_{\tilde{Z}'}$ of $\tilde{D}'$ is flat over Spec W_3. Then the lifting $\tilde{\phi}'$ of $\tilde{\phi}$ (ϕ being the defining equation of Z_i) is a defining equation of $\tilde{Z}'$. We conclude as in of [EV92, Lemma 8.14] that $\tilde{Z}' = \sum \tilde{Z}'_j$ is a lifting over W_3 of $\tilde{Z}$ and we can continue like this for every step.

Since all local liftings can be obtained this way, we can glue and obtain a map $res \in H^0_{crys}(Z) = \oplus W(k)[Z_i]$.

For all this, we have worked with a smooth divisor, namely with the extension of the isocrystal on $X \setminus \text{codim}\, 2$. By [Del70, II 5.8] , and as remarked in [Ked07] we know that if a line bundle with log connection extends to $X \setminus \text{codim}\, 2$ then it extends to the whole of X.

References

[ABV05] F. Andreatta and L. Barbieri-Viale, *Crystalline realizations of 1-motives*, Math. Ann. **331**(2005), no. 1, 111–172.

[BO78] P. Berthelot and A. Ogus, *Notes on crystalline cohomology,* Princeton University Press, 1978.

[BO83] P. Berthelot and A. Ogus, *F-Isocrystals and De Rham Cohomology. I,* Invent. math. **72** (1983), 159–199.

[BI70] P. Berthelot and L. Illusie, *Classes de Chern en cohomologie cristalline*, Comptes Rendus Acad. Sciences Paris **270**(1970), 1695–1697 and 1750–1752.

[Del70] P. Deligne, *Equations Differéntielles à Points Singuliers Réguliers*, Lecture Notes in Math.,**163**, Springer-Verlag, 1970.

[EV92] H. Esnault and E. Viehweg *Lectures on Vanishing Theorems*, Lecture Notes, Birkhäuser, DMV Seminar 20 (1992) 164 pages.

[ES16] H. Esnault and A. Shiho, *Convergent isocrystals on simply connected varieties*, preprint 2016, `www.mi.fu-berlin.de/users/esnault/helene_publ.html`(116).

[ES15] H. Esnault and A. Shiho, *Chern classes of crystals*, preprint 2015, `www.mi.fu-berlin.de/users/esnault/helene_publ.html`(117).

[Esn16] H. Esnault, *Some fundamental groups in arithmetic geometry*, preprint 2016, `www.mi.fu-berlin.de/users/esnault/helene_publ.html`(120).

[EM10] H. Esnault and V. Mehta, *Simply connected projective manifolds in characteristic $p > 0$ have no nontrivial stratified bundles,* Invent. math. **181** (2010), 449–465. (Erratum available at `www.mi.fu-berlin.de/users/esnault/helene_publ.html, 95b` (2013)).

[Gie75] D. Gieseker, *Flat vector bundles and the fundamental group in non-zero characteristics,* Ann. Sc. Norm. Super. Pisa, 4 Sér. **2** (1) (1975), 1–31.

[Gro70] A. Grothendieck, *Représentations linéaires et compactifications profinies des groupes discrets,* Manuscr. Math. **2** (1970), 375–396.

[HyKa] H. Kato, *Semi-stable reduction and crystalline cohomology with logarithmic poles*, Astérisque **223** (1993), 221-268.

[Ill79] L. Illusie, *Complexe de de Rham-Witt et cohomologie cristalline*, Publ. Sc. É.N.S. **12** 4ième série, (1979), 501–661.

[Ka89] K. Kato, *Logarithmic structures of Fontaine-Illusie*, in Algebraic Analysis, Geometry and Number Theory (1989), 191-224.

[Ked07] K. S. Kedlaya, *Semistable reduction for overconvergent F-isocrystals, I: Unipotence and logarithmic extensions*, Compositio Math. **143**(2007), 1164–1212.

[Kin13] L. Kindler, *Evidence for a generalisation of Gieseker's conjecture on stratified bundles in positive characteristic*, Documenta Math. **18** (2013), 1215-1242.

[Mal40] A. Malčev, *On isomorphic matrix representations of infinite groups,* Mat. Sb. N.S. **8**(50) (1940), 405–422.

[Mum70] D. Mumford, *Abelian varieties,* Oxford University Press, London, 1970.

[NaSh08] Y.Nakkajima and A. Shiho, *Weight-filtered convergent complex*, Lecture Notes in Mathematics, Springer Berlin Heidelberg (2008).

[Ogu94] A. Ogus, *F-crystals, Griffiths transversality, and the Hodge decomposition*, Astérisque **221**(1994).

[Pet01] D. Petrequin, *Classes de Chern en cohomologie rigide*, arXiv:math/0103117, 2001.

[Sh00] A. Shiho, *Crystalline Fundamental Groups I Isocrystals on Log Crystalline Site and Log Convergent Site*, J. Math. Sci. Univ. Tokyo, **7**(2000), 509–656.

[Sh02] A. Shiho, *Crystalline fundamental groups II — Log convergent cohomology and rigid cohomology*. J. Math. Sci. Univ. Tokyo, **9**(2002), 1–163.